

# The Inverse 1-Median Problem in $\mathbb{R}^d$ with the Chebyshev-Norm <sup>★</sup>

Johannes Hatzl

*Institute of Optimization and Discrete Mathematics  
Graz University of Technology  
Steyrergasse 30, 8010 Graz, Austria*

*Key words:* 1-median problem, inverse location problem, fractional  $b$ -matching problem, parametric flow problem

---

## 1 Introduction

This paper focuses on the weighted 1-median problem in  $\mathbb{R}^d$  where the distance of two points is measured by the Chebyshev-norm. So far this problem is only well understood for  $d = 2$ . In this case, a linear time algorithm is given in Hamacher [2]. In this note, we give the first combinatorial algorithm for  $d \geq 3$ . Furthermore, we discuss an optimality criterion for the  $d$ -dimensional case which is based on linear programming. Using this optimality criterion we are able to solve the inverse location problem. In the inverse problem the facility is already given and the task is to modify the weights of the points at minimum cost such that the given facility is a 1-median with respect to the new weights.

## 2 The 1-median problem

The 1-median problem discussed in this note is defined as follows: Given  $n$  points  $P_1, \dots, P_n$  with  $P_i = (x_1^i, \dots, x_d^i) \in \mathbb{R}^d$  for  $i = 1, \dots, n$  and associated non-negative weights  $w_i$  the task is to find a point  $P^* = (x_1^*, \dots, x_d^*) \in \mathbb{R}^d$

---

<sup>★</sup> This research has been supported by the Austrian Science Fund (FWF) Project P18918-N18

*Email address:* [hatzl@opt.math.tugraz.at](mailto:hatzl@opt.math.tugraz.at) (Johannes Hatzl).

such that

$$\sum_{i=1}^n w_i \|P_i - P\|_\infty \geq \sum_{i=1}^n w_i \|P_i - P^*\|_\infty$$

for all  $P \in \mathbb{R}^d$ . Note that using some straightforward techniques the problem  $\min_{P=(y_1, \dots, y_d) \in \mathbb{R}^d} \sum_{i=1}^n w_i \|P_i - P\|_\infty$  can be written as a linear programming problem in the following form:

$$\min \quad \sum_{i=1}^n w_i z_i \quad (1)$$

$$\text{s.t.} \quad z_i + y_j \geq x_j^i \quad i = 1, \dots, n, \quad j = 1, \dots, d \quad (2)$$

$$z_i - y_j \geq -x_j^i \quad i = 1, \dots, n, \quad j = 1, \dots, d \quad (3)$$

$$y_j \in \mathbb{R}, \quad z_i \in \mathbb{R} \quad j = 1, \dots, d, \quad i = 1, \dots, n. \quad (4)$$

Due to the fact that the variables  $y_j$  do not appear in the objective function, we use the well known Fourier-Motzkin elimination to get rid of these variables in the constraints. We obtain the following equivalent problem

$$\begin{aligned} \min \quad & \sum_{i=1}^n w_i z_i \\ \text{s.t.} \quad & z_i + z_k \geq d^{ik} \quad i = 1, \dots, n \quad k = 1, \dots, n \\ & z_i \geq 0 \quad i = 1, \dots, n \end{aligned}$$

where  $d^{ik} := \max_j |x_j^i - x_j^k|$ . It is easy to see that the corresponding dual problem is the linear relaxation of the maximum-weight- $b$ -matching problem on a complete graph. It is shown in Antsee [1] that this graphtheoretical problem can be solved by a min-cost-flow problem in a bipartite graph.

### 3 The Inverse Problem

An instance of the inverse problem is given by a set of  $n$  points  $P_1, \dots, P_n \in \mathbb{R}^d$  with corresponding non-negative weights  $w_i \geq 0$  and a point  $P_0$  (which may coincide with a given point). The task is to find new weights  $\tilde{w}_i \geq 0$  such that  $P_0$  is a 1-median with respect to  $\tilde{w}_i$  and  $\|w - \tilde{w}\|_1$  is minimized. The problem can be formulated in a compact form as follows:

$$\begin{aligned} \min \quad & \sum_{i=1}^n |w_i - \tilde{w}_i| \\ \text{s.t.} \quad & \sum_{i=1}^n \tilde{w}_i \|P_i - P\|_\infty \geq \sum_{i=1}^n \tilde{w}_i \|P_i - P_0\|_\infty \quad \forall P \in \mathbb{R}^d \\ & \tilde{w}_i \geq 0 \quad i = 1, \dots, n. \end{aligned}$$

Before we give a combinatorial algorithm we state the following lemma.

**Lemma 1** *There exists an optimal solution  $w^*$  of the inverse location problem such that  $w_i \geq w_i^*$  holds for all  $i = 1, \dots, n$ .*

Let us consider the dual linear programming problem of (1)—(4). We introduce non-negative dual variables  $u_{i,j}$  for the constraints (2) and non-negative dual variables  $v_{i,j}$  for the constraints (3) and obtain

$$\begin{aligned} \max \quad & \sum_{i=1}^n \sum_{j=1}^d x_j^i (u_{i,j} - v_{i,j}) \\ \text{s.t.} \quad & \sum_{j=1}^d (u_{i,j} + v_{i,j}) = w_i \quad i = 1, \dots, n \\ & \sum_{i=1}^n u_{i,j} = \sum_{i=1}^n v_{i,j} \quad j = 1, \dots, d \\ & u_{i,j} \geq 0, \quad v_{i,j} \geq 0 \quad i = 1, \dots, n, \quad j = 1, \dots, d. \end{aligned}$$

In order to interpret the dual problem let us construct the following flow problem: Consider the bipartite graph  $G = (V_1 \cup V_2, E)$  where the set  $V_1$  consists of  $n$  vertices, one for each point  $P_i$  of the 1-median problem. The set  $V_2$  has exactly  $2d$  vertices representing the closed cones

$$Q_j^{\geq} := \{x \in \mathbb{R}^d : x_j \geq 0 \text{ and } |x_j| \geq |x_k| \forall k = 1, \dots, d\}$$

and

$$Q_j^{\leq} := \{x \in \mathbb{R}^d : x_j \leq 0 \text{ and } |x_j| \geq |x_k| \forall k = 1, \dots, d\}$$

for  $j = 1, \dots, d$ . We have an edge  $(P_i, Q_j^{\sim})$  if  $P_i$  is in the cone  $Q_j^{\sim}$  ( $\sim \in \{\leq, \geq\}$ ). Moreover, we set the capacity  $u(e)$  of the edges of the bipartite graph to infinity. Furthermore, we add a source  $s$  and the edges  $(s, P_i)$  for all  $i = 1, \dots, n$  with  $u(s, P_i) = w_i$ . Finally, we introduce a sink  $t$  and an edge from each vertex in  $V_2$  to  $t$  with infinite capacity. We denote this network by  $I(P_1, \dots, P_n, w)$ . Furthermore, a flow in  $I(P_1, \dots, P_n, w)$  is called perfect if the flow on the edges  $(Q_j^{\geq}, t)$  and  $(Q_j^{\leq}, t)$  is equal for all  $j$ . The value of a flow is denoted by  $v(f)$ . Now we can state the following theorem.

**Theorem 2** *Suppose we are given  $n$  points  $P_1, \dots, P_n \in \mathbb{R}^d$  with non-negative weights  $w_i \geq 0$ . Then, the origin  $P^* = (0, \dots, 0)$  is an optimal solution of the 1-median problem if and only if there exists a perfect flow  $f$  in  $I(P_1, \dots, P_n, w)$  such that  $v(f) = \sum_{i=1}^n w_i$ .*

**Example 3** *Suppose we are given the following points:  $P_1 = (-1, 3)$ ,  $P_2 = (2, 2)$ ,  $P_3 = (4, -1)$ ,  $P_4 = (2, -2)$ ,  $P_5 = (0, -2)$  and  $P_6 = (-4, 1)$  with the weights  $w_1 = 3$ ,  $w_2 = 1$ ,  $w_3 = 1$ ,  $w_4 = 2$ ,  $w_5 = 2$  and  $w_6 = 3$ . Then, the corresponding instance  $I(P_1, \dots, P_n, w)$  of the balancing flow problem admits a perfect flow  $f$  with  $v(f) = \sum_{i=1}^n w_i$  (see Figure 1). Thus, we can conclude that the origin is a 1-median.*




Fig. 1. On the left hand side we are given an instance of the 1-median problem where the cone  $Q_1^>$  is highlighted. The right hand side shows the graph of the balancing flow problem (without the supersink  $t$ ). On the edges a perfect flow is given.

It is easy to see that the inverse problem can be solved by the following algorithm:

- Algorithm 1** *Step 1: Construct the instance  $I(P_1, \dots, P_n, w)$*   
*Step 2: Find a perfect flow  $f$  in  $I(P_1, \dots, P_n, w)$  that maximizes  $v(f)$*   
*Step 3: The new weights are given by  $w_i^* = f(s, P_i)$*

The main step is obviously the computation of a maximum perfect flow in Step 2. The maximum perfect flow problem can be reformulated as a parametric flow problem, where capacities on the edges  $(Q_j^>, t)$  and  $(Q_j^<, t)$  are given by a parameter  $\lambda_j$ . If we maximize

$$2 \sum_{j=1}^d \lambda_j$$

such that there exists a flow  $f$  that saturates all the edges entering the supersink  $t$ , the flow  $f$  is a maximum perfect flow. This maximization problem is closely related to parametric flow problems discussed in McCormick [3] and can indeed be solved in polynomial time.

## References

[1] R. Antsee, A polynomial algorithm for  $b$ -matchings: an alternative approach. *Inform. Process. Lett.* (1987), 153–157.  
 [2] H. Hamacher, *Mathematische Lösungsverfahren für planare Standortprobleme*. Braunschweig/Wiesbaden: Vieweg (1995).  
 [3] T. McCormick, Fast algorithms for parametric scheduling come from extensions to parametric maximum flow. In *STOC '96: Proceedings of the twenty-eighth annual ACM symposium on Theory of computing* (1996), 319–328.