

The game chromatic number of 1-caterpillars

Adrien Guignard¹

*Université de Bordeaux
LaBRI UMR 5800
351, cours de la Libération
F-33405 Talence Cedex, France*

Abstract

The game chromatic number of a graph is defined using a two players game. In 1993, Faigle et al. proved that the game chromatic number of trees is at most four. In this paper we investigate the problem of characterizing those trees with game chromatic number three, and settle this problem for 1-caterpillars.

Keywords: game chromatic number, caterpillar, tree, leaf.

1 Introduction

The game chromatic number of a graph G , denoted by $\chi_g(G)$, is defined through a *coloring game* with two players Alice and Bob and a set of k colors. Each move by either player consists of coloring an uncolored node of G with a color i of the set. Adjacent vertices must be colored by distinct colors. The game ends if no more vertices can be colored. Alice wins the game if all vertices are colored. Otherwise, Bob wins.

The game chromatic number $\chi_g(G)$ is the least number of colors for which Alice has a winning strategy in this game. This parameter was introduced by Bodlaender [1] (see also [3] for a recent survey). Since then the problem has attracted considerable attention and has been studied for various classes of graphs [4][5]. For instance, it is proved by Zhu [6] that if P is a planar graph then $\chi_g(P) \leq 17$. Faigle et al. [2] proved that the game chromatic number of every tree is at most four. A natural question in this framework is to characterize the trees with given game chromatic number k , for $1 \leq k \leq 4$. Since the answer is obvious for $k = 1, 2$, our aim is to characterize the set of trees with game chromatic number three. The general characterization seems to be a difficult problem. Therefore, we restrict ourselves to the set of caterpillars and settle the problem for 1-caterpillars.

¹ Email:guignard@labri.fr

2 Definitions and properties

A tree is called a *caterpillar* if a path remains after the removal of all its leaves. This path is called the *spine* of the caterpillar. A *1-caterpillar* is a caterpillar such that every vertex of the spine is a neighbor of exactly one leaf, except for the two extremities of the spine that have two leaves.

Since the game chromatic number of a caterpillar C is at most 4, we will play with three colors and determine whether Alice can complete the coloration of C or not. If it is possible, we will say that Alice wins and otherwise that Bob wins. We denote by $[C, c]$ a caterpillar C equipped with a partial coloring c of its vertices. Such a caterpillar will be simply denoted by C whenever the partial coloring c is clear from the context. To compute the game chromatic number of caterpillars, we have to know for any $[C, c]$ not only who wins if Alice begins, but also if Bob begins.

We thus define the *outcome* of a partially colored caterpillar C , denoted by $o(C)$, as follows:

- (i) $o(C) = \mathcal{B}$ if **Bob** wins whoever starts the game;
- (ii) $o(C) = \mathcal{P}$ if the next player loses (so the **Previous** one wins);
- (iii) $o(C) = \mathcal{N}$ if the **Next** player wins;
- (iv) $o(C) = \mathcal{A}$ if **Alice** wins whoever starts the game.

We denote by $Opt(C)$ the set of options of C , that is the set of partially colored caterpillars that can be obtained from C after one move. If C has an option with outcome \mathcal{X} , we say that C has an \mathcal{X} -*option*. We extend the definition of outcome to a set of caterpillars \mathcal{C} : $o(\mathcal{C}) = \{o(C), C \in \mathcal{C}\}$.

Proposition 2.1 *Let C be a not totally colored caterpillar.*

- (i) $o(C) = \mathcal{B} \Leftrightarrow o(Opt(C)) \in \{\{\mathcal{B}\}, \{\mathcal{N}, \mathcal{B}\}\}$;
- (ii) $o(C) = \mathcal{P} \Leftrightarrow o(Opt(C)) = \{\mathcal{N}\}$;
- (iii) $o(C) = \mathcal{N} \Leftrightarrow o(Opt(C))$ contains \mathcal{P} , or contains \mathcal{A} and \mathcal{B} ;
- (iv) $o(C) = \mathcal{A} \Leftrightarrow o(Opt(C)) \in \{\{\mathcal{A}\}, \{\mathcal{A}, \mathcal{N}\}\}$.

Since we will need to consider outcomes of disjoint unions of caterpillars, we need some properties to compute $o(C_1 \cup C_2)$ according to $o(C_1)$ and $o(C_2)$ (where $C_1 \cup C_2$ denotes the disjoint union of C_1 and C_2).

We call the *signed outcome* of C the outcome \mathcal{X} of C signed by the parity of the number of uncolored nodes of C , denoted by $o'(C) = \mathcal{X}_0$ or \mathcal{X}_1 .

Proposition 2.2 *Firstly, if $o(C_1) = \mathcal{B}$ or $o(C_2) = \mathcal{B}$ then $o(C_1 \cup C_2) = \mathcal{B}$. Otherwise, we define an addition function of signed outcomes, denoted by " \oplus ", that satisfies $o'(C_1 \cup C_2) = o'(C_1) \oplus o'(C_2)$, and is given by the following table:*

\oplus	\mathcal{P}_0	\mathcal{P}_1	\mathcal{N}_0	\mathcal{N}_1	\mathcal{A}_0	\mathcal{A}_1
\mathcal{P}_0	\mathcal{P}_0	\mathcal{B}_1	\mathcal{B}_0	\mathcal{N}_1	\mathcal{P}_0	\mathcal{N}_1
\mathcal{P}_1	\mathcal{B}_1	\mathcal{B}_0	\mathcal{B}_1	\mathcal{N}_0	\mathcal{P}_1	\mathcal{N}_0
\mathcal{N}_0	\mathcal{B}_0	\mathcal{B}_1	\mathcal{B}_0	\mathcal{B}_1	\mathcal{N}_0	\mathcal{N}_1
\mathcal{N}_1	\mathcal{N}_1	\mathcal{N}_0	\mathcal{B}_1	\mathcal{B}_0	\mathcal{N}_1	\mathcal{N}_0
\mathcal{A}_0	\mathcal{P}_0	\mathcal{P}_1	\mathcal{N}_0	\mathcal{N}_1	\mathcal{A}_0	\mathcal{A}_1
\mathcal{A}_1	\mathcal{N}_1	\mathcal{N}_0	\mathcal{N}_1	\mathcal{N}_0	\mathcal{A}_1	\mathcal{A}_0

We note \preceq the relation on the set of outcomes $\{\mathcal{B}, \mathcal{P}, \mathcal{N}, \mathcal{A}\}$ such that:

- (i) $\mathcal{A} \preceq \mathcal{N} \preceq \mathcal{B}$
- (ii) \mathcal{P} and every another outcome \mathcal{X} are incomparables.


If T and U are two partially colored caterpillars, we say that T is a subgraph of U and note $T \subseteq U$ if and only if $V(T) \subseteq V(U)$, $E(T) \subseteq E(U)$ and $\forall v \in V(T)$, $c_T(v) = c_U(v)$ (where $c_T(v)$ is the color of v in T).

Proposition 2.3 *Let T and U be two partially colored caterpillars. If $T \subseteq U$ then $o(T) \preceq o(U)$*

Let C be a partially colored caterpillar. Observe that if C has an uncolored vertex v having three neighbours colored with distinct colors, then the outcome is \mathcal{B} (since v cannot be colored). Similarly, if C has a colored node v with degree $k \geq 2$, the forest C' obtained from C by splitting v into k colored leaves with the same color than v , each linked to a neighbour of v (thus creating k connected components) is equivalent to C (we mean $o'(C') = o'(C)$). Finally, if C has an uncolored node v with two leaves colored with the same color, the caterpillar C' obtained from C by deleting one of these two leaves is equivalent to C .

3 The family of 1-caterpillars

Let C be a partially colored 1-caterpillar whose spine $s_1 s_2 \dots s_\ell$ contains no colored vertices. Moreover let s_0 (resp. $s_{\ell+1}$) be one of the two leaves connected to s_1 (resp. s_ℓ). The leaves connected to s_1 or s_ℓ are the *ends* of C . We associate with C a word $w(C) = w_0 \dots w_{\ell+1}$ on the alphabet $\{z, 1, 2, 3\}$ defined as follows: w_0 is the color of s_0 if it is colored or z otherwise (the same is true of $w_{\ell+1}$), and for every i , $1 \leq i \leq \ell$, w_i is the color of the leaf connected to s_i , or z if this leaf is not colored. For instance, the following caterpillar is associated with the word $1zz2z2$.



Using properties of outcomes and subgraphs, we prove the following results.

Theorem 3.1 *Let C be a 1-caterpillar with $w(C) = az^nb$ and $a, b \in \{1, 2, 3\}$. The outcome of C is given by the following table:*

n	1	2	3	4	5	6	7	8	9
$o(C)$	\mathcal{AN}	\mathcal{A}	\mathcal{N}	\mathcal{A}	\mathcal{N}	\mathcal{A}	\mathcal{N}	\mathcal{AN}	\mathcal{N}
n	10	11	12	13	14	15	16	17	≥ 18
$o(C)$	\mathcal{N}	\mathcal{N}	\mathcal{N}	\mathcal{N}	\mathcal{NB}	\mathcal{N}	\mathcal{B}	\mathcal{NB}	\mathcal{B}

where \mathcal{XY} stands for \mathcal{X} if $a = b$ and \mathcal{Y} otherwise.

The proof relies on several lemmas which consider the outcomes of some 1-caterpillars with particular partial colorings. We proceed by studying a family \mathcal{F} of caterpillars (for instance $az^{2n}bb$ with $n \geq 0$) and compute the value n defined as the smallest size of a caterpillar with outcome \mathcal{N} (resp. \mathcal{B}), so that every smaller caterpillar in \mathcal{F} has outcome \mathcal{A} (resp. \mathcal{A} or \mathcal{N}). We also have to check that no 1-caterpillar of this family has outcome \mathcal{P} .

Theorem 3.2 *Let C an uncolored 1-caterpillar with n nodes of degree 3.*

- (i) *If $n \geq 28$ then $o(C) = \mathcal{B}$*
- (ii) *If $23 \leq n \leq 27$ then $o(C) = \mathcal{N}$*
- (iii) *Otherwise $o(C) = \mathcal{A}$*

References

- [1] H. L. Boadlaender, *On the complexity of some coloring games*, Int. J. Found Comput. Sci. **2**(2) (1991), 133–147.
- [2] U. Faigle, U. Kern, H. A. Kierstead and W. T. Trotter, *On the game chromatic number of some classes of graphs*, Ars Combin **35** (1993), 143–150.
- [3] T. Bartnicky, J. Grytczuk, H. A. Kierstead and X. Zhu, *The map coloring game*, Am. Math. Monthly, **114** (2007), 793–803.
- [4] H. A. Kierstead and W. T. Trotter, *Planar graph coloring with an uncooperative partner*, J. Graph Theory **18**(6) (1994), 569–584.

- [5] D. Guan and X. Zhu, *The game chromatic number of outerplanar graphs*, J. Graph Theory **30**(1999), 67–70.
- [6] X. Zhu, *Refined activation strategy for the marking game*, J. Combin. Theory Ser. B **98**(1) (2008), 1–18